

SUMMARY OF RECOMMENDED PERIODS OF ABSENCE FOR COMMUNICABLE DISEASES

ADULTS

Health Protection Team, Lothian NHS Board, Waverley Gate, 2-4 Waterloo Place, Edinburgh, EH1 3EG
Enquiries 0131 465 5420/5422

**LOTHIAN NHS BOARD
HEALTH PROTECTION TEAM
Recommended Periods of Absence for Communicable Disease - Adults**

Introduction

In accordance with national guidance, this document provides a summary of the recommended periods of absence for adults who are, or are thought to be, suffering from an infection which may spread to others.

The key to prevention and control of spread of infection is maintaining high standards of hygiene at all times. In particular:

Hands should be washed **after**:

- Using or cleaning the toilet
- Immediately after handling raw meat/poultry
- Preparing food in general
- Blowing your nose or covering a sneeze or cough
- Smoking, as fingers will come in contact with the mouth and nose
- Touching pets or other animals

Hands should be washed **before**:

- Preparing food or drink
- Eating food

High standards of hygiene should also be applied to:

- Food hygiene
- Environmental cleaning
- Disposal of waste.

For further information and advice on infectious diseases and the control of infection, contact the Health Protection Team, Lothian NHS Board.

Tel: 0131 465 5420/5422

**LOTHIAN NHS BOARD
HEALTH PROTECTION TEAM
Recommended Periods of Absence for Communicable Disease - Adults**

Key for recommended period of absence:

Groups that pose a higher than normal risk of spreading infection.

Group A	Any person of doubtful hygiene or with unsatisfactory toilet, hand washing or hand drying facilities at home, work or school.
Group B	Children who attend pre-school groups or nursery
Group C	People whose work involves preparing or serving unwrapped foods not subjected to further heating/cooking.
Group D	Health or Social Care staff who have direct contact with highly susceptible patients or persons in whom an infection would have particularly serious consequences.

An outbreak is usually defined by the presence of 2 or more cases with the same suspected or confirmed disease. The size of the total population also needs to be taken into account but if you are unsure then please call HPT to discuss.

**LOTHIAN NHS BOARD
HEALTH PROTECTION TEAM**
Recommended Periods of Absence for Communicable Disease - Adults

Disease/ Causative Organism	Typical Incubation Period	Route of Infection	Risk of person to person spread	Recommended Period of Absence		Action
				Cases	Contacts	
Campylobacter	3-4 Days (Can be 1-10 days).	Food borne. Contaminated food and water. Contact with infected animals.	Low risk of transmission from person to person can occur, especially where there is poor hygiene practice.	Until clinically recovered and diarrhoea has ceased for 48hrs.	None	Practice good hygiene, specifically hand hygiene.
Chickenpox (Varicella Zoster)	15-18 days (Can be 10-21 days).	Direct person to person contact – airborne and droplet spread.	High risk of transmission from 2 days before rash onset until all the lesions have crusted.	Until vesicles become dry (approx 7 days) but a minimum of 5 days after rash onset.	Healthcare workers without a history of chickenpox may be excluded by HPT.	Pregnant women and the immuno-compromised who are contacts of cases should seek medical advice as soon as possible.
Clostridium Difficile	Variable. Often triggered by antibiotic use and can start a few days or months after antibiotic course.	Contact with an infected person or contaminated environment or objects.	Those most at risk of transmission are elderly, people currently or recently taken antibiotics, been in hospital or are immunosuppressed.	Until clinically recovered and diarrhoea has ceased for 48hrs.	None	Practice good hygiene, specifically environmental and hand hygiene.
Colds	12hours to 5 days (Commonly 48 hours).	Respiratory droplet. Contact with secretions.	High risk of transmission during active infection.	None.	None.	Practice good hygiene. Practice good cough etiquette
Cold sores	2-12 days.	Direct contact with oral	High risk of transmission	None.	None.	Practice good hygiene.

**LOTHIAN NHS BOARD
HEALTH PROTECTION TEAM**

Recommended Periods of Absence for Communicable Disease - Adults

Disease/ Causative Organism	Typical Incubation Period	Route of Infection	Risk of person to person spread	Recommended Period of Absence		Action
				Cases	Contacts	
(Herpes Simplex)		secretions or direct contact with lesion.	until lesion crusted.			Health education. Avoid kissing and contact with sores. Mild self limiting.
Conjunctivitis	1-3 days.	Contact with discharges.	High risk of transmission whilst symptomatic.	If unwell and a serious infection stay off until eye no longer inflamed and infected.	None.	Practice good hygiene. If outbreak or cluster inform HPT.
Cryptosporidiosis	7-10 days (Can be 1-21 days average 7 days).	Faecal-oral. Waterborne. Contact with animal faeces.	High risk of transmission from person to person.	Until clinically recovered and diarrhoea has ceased for 48hrs. Avoid swimming until 2 weeks after symptoms cease.	None.	Practice good hygiene. Follow up by HPT/Environmental health.
Cytomegalovirus (CMV)	Variable. 3-8 weeks but can be up to 12.	Intimate exposure. Contact with infected tissue or fluids (e.g. body fluids/blood).	High risk of transmission through intimate contact with fluids, whilst organism present. Carriage may persist for many months.	None.	None.	Practice good hygiene.
Diarrhoea	Dependent on causative organism.	Often food or waterborne or due to poor hygiene. Can be faecal-oral. Some viruses may be airborne.	High risk of transmission whilst symptomatic, though dependent on cause.	Until clinically recovered and diarrhoea has ceased for 48hrs. (If cause known refer to disease).	None (If cause known refer to disease).	Practice good hygiene (If cause known refer to disease).

**LOTHIAN NHS BOARD
HEALTH PROTECTION TEAM**

Recommended Periods of Absence for Communicable Disease - Adults

Disease/ Causative Organism	Typical Incubation Period	Route of Infection	Risk of person to person spread	Recommended Period of Absence		Action
				Cases	Contacts	
Diphtheria (very rare in UK)	2-5 days but may be longer.	Contact with discharge from lesions. Airborne droplet spread. Direct contact with respiratory discharges. Infected animals or unpasteurised dairy products.	Not highly infectious. Prolonged close contact is normally required for transmission. Cases remain infectious for up to 4 weeks after symptom onset or after 3 days of appropriate antibiotics.	Until clinically recovered and bacteriological specimens are clear. Always consult with local health protection team as exclusion will apply.	Household contacts should be excluded until specimens are clear.	Notifiable. Investigation by HPT Preventable by vaccination. Contact tracing will be required.
Dysentery (Bacillary) = <i>Shigella flexneri</i> <i>Shigella boydii</i> <i>Shigella dysenteriae</i> <i>*Shigella sonnei</i>	1-3 days (Can be 8hrs to 7 days).	Faecal-oral. Food borne. Occasionally waterborne	High risk of transmission from person to person especially whilst cases are symptomatic.	=2 negative stool specimens for groups A, B, C and D taken at least 48 hours apart. * 48 hours symptom free.	None. Unless contacts are symptomatic	Notifiable Practice good hygiene. Follow up by EHO/HPT.
E.coli O157 (VTEC)	2-7 days (Can be 1-14).	Faecal-oral. Food borne. Contact with animal faeces.	Very high risk of transmission from person to person. Cases can still pass on infection once asymptomatic and contacts can carry E-coli O157 with no symptoms and pass it on.	2 negative stool specimens for groups A, B, C and D taken 24 hours apart. Otherwise until clinically recovered and diarrhoea has ceased for 48 hrs.	2 negative stool specimens for groups A, B, C and D taken 24 hours apart. Otherwise 48 hrs symptom free.	Notifiable. Follow up by HPT/EHO. Practice good hygiene
E.coli in urine	Unknown.	Spread of the bacteria from the gut to the urinary system.	Low risk of transmission from person to person.	Until clinically recovered, usually 2-4 days. May require short course of	None	Practice good hygiene.

**LOTHIAN NHS BOARD
HEALTH PROTECTION TEAM**

Recommended Periods of Absence for Communicable Disease - Adults

Disease/ Causative Organism	Typical Incubation Period	Route of Infection	Risk of person to person spread	Recommended Period of Absence		Action
				Cases	Contacts	
				antibiotics.		
Fifth Disease (Parvovirus B19 or 'slapped-cheek' disease)	13-18 days (4-20 days).	Through contact with respiratory secretions.	High risk of transmission 7 days before rash appears until one day after onset of rash. Most adults are immune to parvovirus.	Until clinically recovered.	None.	Pregnant women, immunocompromised and people with haemolytic anaemia should avoid contact with known cases.
Food Poisoning	Dependent on causative organism.	Food borne.	Variable. Usually low risk of transmission if asymptomatic.	Until clinically recovered and diarrhoea has ceased for 48hrs. (If cause known refer to disease).	None. (If cause known refer to disease).	Practice good hygiene.
German Measles (Rubella)	14-17 days (Can be 14-21 days).	Droplet spread or direct contact with secretions.	High risk of transmission from 1 week before onset of rash to 4 days after.	Until clinically recovered but <u>at least</u> 4 days after onset of rash.	If contacts work with vulnerable groups or in large institutions they may be excluded by HPT.	Notifiable. Follow up of cases by HPT. Pregnant women should consult their GP or midwife if exposed. Children should be immunised with MMR.
Giardiasis	7-10 days (Can be 5-28 days).	Waterborne. Faecal-oral.	High risk of transmission whilst organism present in stool.	Until clinically recovered and diarrhoea has ceased for 48hrs	None.	Notifiable. Follow up by HPT/EHO. Practice good hygiene.
Glandular Fever (Infectious Mononucleosis)	4-6 weeks.	Close contact with pharyngeal secretions (e.g. kissing). Indirectly on hands.	Carriage may be prolonged with high risk of transmission.	Until clinically recovered.	None.	

**LOTHIAN NHS BOARD
HEALTH PROTECTION TEAM**

Recommended Periods of Absence for Communicable Disease - Adults

Disease/ Causative Organism	Typical Incubation Period	Route of Infection	Risk of person to person spread	Recommended Period of Absence		Action
				Cases	Contacts	
Group A streptococcal infection	1-4 days for acute infection, 2-3 weeks for untreated sore throat	Person to person	Low risk of transmission	Until clinically recovered and 24 hours after appropriate antibiotics	None unless symptomatic. Information will be given to close contacts	HPT will investigate cases of invasive group A strep
Haemophilus Influenzae B (HIB)	Unknown but probably 2-4 days.	Respiratory droplet or contact with secretions.	High risk of transmission whilst symptomatic and/or whilst organism is present in nasopharynx. Non infectious after 48 hours of appropriate antibiotic treatment.	Until clinically recovered but at least 48 hours after commencing treatment.	None.	Investigation by HPT. Children should have been immunised pre-school.
Hand, Foot & Mouth Disease	3-5 days.	Direct contact with faeces, blisters and respiratory droplets (aerosol droplet spread).	High risk of transmission during acute stage of illness (occasionally longer as virus can persist in faeces for several weeks).	When clinically recovered.	None.	Report outbreaks (2 or more) to HPT. Practice good hygiene.
Head Lice	Head lice mature in 6-12 days and live for about 20 days.	Direct head to head contact.	High risk of transmission until adequately treated.	Until treated. (After first treatment and no visible live lice). Treatment only recommended when live lice are seen.	None if asymptomatic.	Health education. Practice good hygiene.
Hepatitis A	28-30 days (Can be 15-50).	Faecal-oral. Waterborne.	High risk of transmission from two weeks before onset of jaundice until one week after jaundice starts.	Until 7 days after onset of jaundice (if present) or other symptoms.	None. (unless they have symptoms suggestive of hepatitis A or are food handlers)	Notifiable. Follow up by HPT/EHO. Practice good hygiene.

**LOTHIAN NHS BOARD
HEALTH PROTECTION TEAM**

Recommended Periods of Absence for Communicable Disease - Adults

Disease/ Causative Organism	Typical Incubation Period	Route of Infection	Risk of person to person spread	Recommended Period of Absence		Action
				Cases	Contacts	
Hepatitis B	2-6 months (Commonly 2-3).	Blood borne. Mother to baby vertical transmission. Sexual transmission. Sharing injecting equipment.	Infectious during incubation period and up to 6 months after acute illness. 10% of cases develop chronic infection and continue to be infectious Low risk of transmission if social contact only.	Until clinically recovered.	None.	Notifiable. Investigation by HPT. Practice good hygiene, with care when dealing with blood and body fluids. Practice safe sex Do not share needles.
Hepatitis C	2 weeks to 6 months (commonly 6-10 weeks)	Blood borne. Sexual transmission. Sharing injecting equipment.	Probably infectious for life. Low risk of transmission if social contact only.	Until clinically recovered.	None.	Notifiable. Practice good hygiene with care when dealing with blood/body fluids.
Hepatitis E	15-64 days Mean value 26-42 days	Faecal-Oral Contaminated food/water	Virus present in faeces during late incubation and have been detected 14 days after onset of jaundice. No evidence of person-person transmission.	Until clinically recovered but all should be reminded of the importance of hand hygiene.	None	Notifiable Investigation by HPT/EHO Hand Hygiene.
HIV	1-3 months for detectable antibodies	Blood borne. Sexual transmission. Sharing injecting equipment. Mother to child vertical transmission.	Infectious for life. Low risk of transmission if social contact only.	None.	None.	Notifiable. Practice good hygiene. take care when dealing with blood/body fluids.
Impetigo	1-10 days.	Direct contact with lesions. Indirect contact with infected items (e.g. towels, clothes).	Highly infectious whilst lesions are present and until they are healed and crusted over.	Until lesions are crusted or healed or 48 hours after commencing appropriate antibiotics.	None.	Report outbreaks to HPT. Practice good hygiene.
Influenza	1-5 days.	Airborne/droplet. Contact with	Highly infectious in the first 3-5 days (up to 10	Until clinically recovered.	None.	Immunisation for at risk groups.

**LOTHIAN NHS BOARD
HEALTH PROTECTION TEAM**

Recommended Periods of Absence for Communicable Disease - Adults

Disease/ Causative Organism	Typical Incubation Period	Route of Infection	Risk of person to person spread	Recommended Period of Absence		Action
				Cases	Contacts	
		respiratory secretions.	days in young children).			Practice good hygiene.
Measles	7-18 days (can be up to 21 days). Rash usually appears 14 days after exposure.	Airborne. Direct contact with respiratory secretions.	Highly infectious from 5 days before onset of rash until 4 days after the rash develops.	4 days from the onset of rash.	Contacts who work with vulnerable people or in large institutions may be excluded by HPT	Notifiable Investigation by HPT Children should be routinely immunised with MMR. Pregnant women should seek advice from their GP/midwife.
Meningococcal Disease/Septicemia	2-10 days. Commonly 3-4.	Direct contact. Contact with respiratory droplets from nose and throat.	Low risk of transmission person to person until 48 hrs of appropriate antibiotic therapy.	Until clinically recovered.	None.	Notifiable. Investigation by HPT. Meningitis ACWY vaccination recommended for 14-18 year olds and University fresher's. Men B vaccine now also part of childhood immunisation schedule.
Mumps	16-18 days (Can be 12-25 days).	Airborne/droplet spread. Direct contact with saliva.	Medium risk of transmission 7 days before onset of symptoms until 9 days after.	Until clinically recovered but no less than 5 days from the onset of symptoms/ swollen glands.	None.	Notifiable. Preventable by vaccination with 2 X MMRs. Inform HPT if outbreak suspected.
Norovirus (Winter vomiting bug)	15-50 hours (can be 4-77 hours)	Faecal-Oral Aerosol transmission	High risk of transmission from person to person and environmental transmission.	Until recovered. 48 hours since symptoms have ceased.	None	Inform HPT if outbreak suspected. Practice good hygiene
Poliomyelitis (very rare in UK)	7-14 days (Can be 3-35 days).	Faecal-oral.	High risk of transmission when virus present in stools and/or nasopharynx.	At the discretion of the duty Consultant in Public Health Medicine (CPHM)	None.	Notifiable. Investigation by HPT. Children should be routinely immunised.
Ringworm	2-6 weeks.	Direct skin to skin	Medium risk of	None, but lesions	Families	Avoid direct contact with

**LOTHIAN NHS BOARD
HEALTH PROTECTION TEAM**

Recommended Periods of Absence for Communicable Disease - Adults

Disease/ Causative Organism	Typical Incubation Period	Route of Infection	Risk of person to person spread	Recommended Period of Absence		Action
				Cases	Contacts	
		contact with infected person or animal. Indirect contact with fomites or environmental surfaces.	transmission whilst infected lesions are present.	should be covered.	should be checked for ringworm.	lesions. Good hygiene practice.
Rubella (see German Measles)						
Salmonella (excluding typhoid and paratyphoid)	12-36 hrs (can be 6hrs to 7 days)	Faecal-oral Contaminated food	High risk of transmission when symptomatic.	None but until clinically recovered and 48hrs after diarrhoea has ceased.	None	Notifiable. Follow up by HPT/EHO. Practice good hygiene
Scabies	2-6 weeks if not previously infected. 1-4 days if re-infected.	Prolonged skin to skin contact. E.g. hand holding.	High risk of transmission until adequately treated.	Until treated. Can return after first treatment.	All household and close contacts may require treatment	Practice good hygiene. Health education. Contact HPT if two or more cases.
Scarlet Fever	1-3 days.	Airborne/droplet. Contact with respiratory secretions. Direct contact with patients or carriers.	Medium risk of transmission whilst organism present in nasopharynx, although minimal risk after 24 hours of appropriate antibiotic treatment.	Until clinically recovered and 24 hours after start of treatment.	None.	None.
Shigella (see Dysentery)						
Shingles (Varicella-Zoster)	Reactivation of Varicella infection	Direct contact with lesions.	Moderate risk of transmitting chickenpox	None if lesions can be covered and are not	None. Can cause	Practice good hygiene. Seek advice from GP or

**LOTHIAN NHS BOARD
HEALTH PROTECTION TEAM**

Recommended Periods of Absence for Communicable Disease - Adults

Disease/ Causative Organism	Typical Incubation Period	Route of Infection	Risk of person to person spread	Recommended Period of Absence		Action
				Cases	Contacts	
virus)	(chickenpox).		in the 7 days after the appearance of lesions.	weeping. Otherwise for 7 days after onset of lesions.	Chicken Pox in those who have not had it	midwife if pregnant/immunocompromised.
Staphylococcus aureus infection (MRSA/MSSA/PVL)	Variable and indefinite	Contact with patients with purulent lesions, or with an asymptomatic carrier. Airborne spread is rare	High for certain groups of patients, and if patients have active infection and there is skin to skin contact, poor hygiene or sharing of personal items.	Special considerations for Health care workers: contact occupational health.	None	HPT will only follow up cases of PVL staph aureus. Hand hygiene is very important. Health education should be provided. Any active lesions should be covered.
Threadworm	Variable- days to weeks.	Faecal oral.	Medium risk of transmission whilst eggs in stool.	None but should be treated properly.	Household contacts should be treated at the same time as the case.	Practice good hygiene. Health education.
Thrush (candidiasis)	Variable. 2-5 days in infants.	Contact with secretions from mouth, skin, vagina and faeces. Vertical mother to child transmission at birth.	High risk of transmission.	None.	None.	Practice good hygiene. Health education.
Toxocariasis	Weeks or months depending on the severity of infection.	Ingestion of eggs from contaminated soil, hands or contact with dogs (especially puppies).	Not spread from person to person.	None.	None.	Practice good hygiene. Health education.
Toxoplasmosis	5-23 days.	Ingestion of eggs from sand boxes/play areas contaminated with cat faeces.	Not spread from person to person.	None.	None.	Practice good hygiene. Health education.

**LOTHIAN NHS BOARD
HEALTH PROTECTION TEAM**

Recommended Periods of Absence for Communicable Disease - Adults

Disease/ Causative Organism	Typical Incubation Period	Route of Infection	Risk of person to person spread	Recommended Period of Absence		Action
				Cases	Contacts	
		Also from rare, undercooked meats.				
Tuberculosis – pulmonary	2-12 weeks (Disease can be “sleeping” for decades).	Airborne/droplet.	Medium to low risk until 2 weeks after treatment. Requires close prolonged contact.	After consultation with specialist physician the TB nurse/CPHM will inform the patient about their return to work.	At the discretion of the CPHM.	Notifiable. Investigation by HPT. At risk children should be vaccinated as babies.
Tuberculosis – non-pulmonary	Indefinite.	Not usually infectious.	Not usually transmitted from person to person.	Until clinically recovered.	None.	Notifiable Investigation by HPT.
Typhoid and Paratyphoid Fever	10-14 days (Can be 1-3 weeks).	Food borne. Waterborne. Faecal/Urine-oral.	High risk of transmission whilst symptomatic.	3 negative stool specimens 48hrs apart if in a risk group (starting at least one week after antibiotic course completed). Otherwise 48 hrs symptom free.	None unless symptomatic in which case should be excluded until symptoms have ceased for at least 48hrs.	Notifiable. Follow up by HPT/EHO. Practice good hygiene.
Viral Gastro- enteritis	Dependant on causative organism.	Faecal-oral. Airborne secondary to environmental contamination. Food borne.	High risk of transmission during acute vomiting and diarrhoea and up to 48 hrs after symptom free.	48hrs after symptoms cease.	None.	Outbreaks followed up by HPT/EHO’s. Practice good hygiene.
Viral Meningitis	Dependant on virus.	Dependant on the virus. Person to person spread usually droplet spread or may be airborne.	Considered to be very unlikely.	Until clinically recovered (usually within a week).	None.	If more than one case then may wish to speak to HPT for advice. Practice good hygiene.
Vomiting	Dependant on causative organism.	Often food or waterborne or due to poor hygiene. Can be faecal-oral.	High risk of transmission whilst symptomatic, though dependent on cause.	Until clinically recovered and symptoms have ceased for 48 hrs.	None.	Outbreaks followed up by HPT/EHO’s. Practice good hygiene.

**LOTHIAN NHS BOARD
HEALTH PROTECTION TEAM**

Recommended Periods of Absence for Communicable Disease - Adults

Disease/ Causative Organism	Typical Incubation Period	Route of Infection	Risk of person to person spread	Recommended Period of Absence		Action
				Cases	Contacts	
		Some viruses may be airborne.				
Warts/Verrucae	2-3 months (But can be 1-20 months).	Direct contact with warts. Verrucas may spread in pools or showers.	Can probably be transmitted at least as long as visible lesions present.	None. Verrucae should be covered in swimming pools, gyms and changing rooms.	None.	Practice good hygiene.
Whooping Cough (pertussis)	7-10 days (Can be 5-21).	Airborne/droplet. Contact with respiratory secretions.	Highly infectious in early stages of illness and up to 3 weeks after onset of cough (rarely 6 weeks).	Until clinically recovered and 3 weeks from onset of cough or 48hrs of appropriate antibiotic therapy.	If symptomatic see case absence.	Notifiable. Investigation by HPT. Children should be routinely immunised pre-school.

Health Protection Team, Lothian NHS Board, Waverley Gate, 2-4 Waterloo Place, Edinburgh, EH1 3EG
Enquiries 0131 465 5420/5422